



RABEJAZE Tablet

Proton Pump Inhibitor for Acid-Related Disorders

Relieve acid reflux. Heal ulcers. Protect your gut.

Product Overview

RABEJAZE is a proton pump inhibitor (PPI) used to treat a variety of acid-related gastrointestinal conditions. It reduces stomach acid production by inhibiting the gastric proton pump, an enzyme responsible for acid secretion. RABEJAZE promotes ulcer healing and helps prevent new ulcers from forming.

Active Ingredient

- Rabeprazole – 20 mg

Key Benefits

- Alleviates heartburn and gastroesophageal reflux disease (GERD)
- Heals erosive esophagitis
- Treats and prevents duodenal and gastric ulcers
- Part of therapy for **Helicobacter pylori** infection (in combination with antibiotics)
- Reduces excessive acid in Zollinger-Ellison syndrome

How It Works

Rabeprazole inhibits the gastric proton pump (H^+/K^+ ATPase), effectively blocking acid production at its source in the stomach. This leads to reduced acidity, faster ulcer healing, and relief from reflux symptoms.

How to Use

- Take 30 minutes before meals
- Swallow whole with water—do not crush or chew
- Follow your doctor's prescribed dose and schedule

Storage

- Store in a cool, dry place away from sunlight
- Keep out of reach of children

Possible Side Effects

- Headache, dizziness
- Nausea, vomiting, diarrhea, constipation
- Flatulence, abdominal discomfort
- Weakness or runny nose
- Dry mouth (rare)

If symptoms persist or worsen, consult your doctor.

Warnings & Precautions

- Do not take if allergic to Rabeprazole or other PPIs
- Use with caution in elderly patients due to fracture risk
- Monitor magnesium and vitamin B12 levels during long-term use
- Inform your doctor if pregnant or breastfeeding
- Regular check-ups may be needed during prolonged therapy

Frequently Asked Questions (FAQs)

1. What is RABEJAZE used for?

It treats GERD, duodenal ulcers, erosive esophagitis, **H. pylori** infections (with antibiotics), and Zollinger-Ellison syndrome.

2. How long can I take RABEJAZE?

Short-term use is preferred unless your doctor advises otherwise. Regular check-ups are important during long-term therapy.

3. Can I stop taking RABEJAZE once I feel better?

No. Complete the course as prescribed. Stopping early may cause symptoms to return.

4. Can RABEJAZE cause diarrhea?

Yes. Stay hydrated and consult your doctor if diarrhea persists or is severe.

5. Can it be taken on an empty stomach?

Yes. Taking it 30 minutes before meals improves its effectiveness.

6. Does it cause vitamin B12 deficiency?

Prolonged use may affect B12 absorption. Speak to your doctor if you experience fatigue, mouth ulcers, or tingling sensations.

7. Is it safe during pregnancy or breastfeeding?

Consult your doctor before use during pregnancy or lactation.

8. Does RABEJAZE increase the risk of bone fractures?

May be yes, especially in elderly patients and those on long-term therapy. Weigh the benefit and risk. Always discuss with your doctor if you have osteoporosis or use corticosteroids.

9. Can it cause dry mouth?

Yes. Drinking water frequently or chewing sugar-free gum may help relieve dryness.

10. What should I do if I miss a dose?

Take it as soon as you remember unless it's close to your next dose. Do not double the dose.

Your Gut, Protected and Healed

RABEJAZE offers fast, effective relief from acid-related conditions—ensuring long-term gut comfort and health.