

OSVYNAC-MR Tablet

Effective Relief from Acute Muscular Pain and Stiffness

Ease discomfort. Improve mobility. Regain function.

Product Overview

OSVYNAC-MR Tablet is a combination medication designed to provide rapid and effective relief from muscular pain. It targets pain, inflammation, and muscle stiffness by inhibiting key chemical mediators in the body. This enhances overall mobility, reduces discomfort, and supports a more active lifestyle.

Active Ingredients & Their Roles

Aceclofenac (100 mg)

- Nonsteroidal anti-inflammatory drug (NSAID)
- Reduces pain and inflammation by blocking COX enzymes

Paracetamol (325 mg)

- Provides additional pain relief
- Helps lower fever

Chlorzoxazone (500 mg)

- Centrally acting muscle relaxant
- Relieves muscle stiffness and spasms

Benefits of OSVYNAC-MR Tablet

- Reduces muscle pain, stiffness, and spasms
- Enhances mobility and physical function
- Improves recovery from muscle injuries, strains, and sprains
- Offers combined anti-inflammatory, analgesic, and muscle-relaxing action

How It Works

- **Aceclofenac** reduces the production of prostaglandins that cause pain and inflammation.
- **Paracetamol** enhances pain relief and lowers fever.

- **Chlorzoxazone** acts on the central nervous system to relieve muscle tension. Together, these ingredients relieve pain, reduce swelling, and restore muscle flexibility.

Uses of OSVYNAC-MR Tablet

- Musculoskeletal pain
- Muscle strains and sprains
- Back pain, neck pain, and shoulder pain
- Post-workout soreness or injury-related stiffness

Directions for Use

- Take exactly as prescribed by your physician
- Preferably consume with or after food to minimize gastric discomfort
- Swallow whole with water — do not chew or crush

Storage Guidelines

- Store in a cool, dry place away from sunlight
- Keep out of reach of children

Common Side Effects

- Nausea
- Vomiting
- Heartburn
- Abdominal discomfort
- Diarrhea
- Loss of appetite
- Drowsiness or fatigue

If any of these symptoms persist, consult your doctor.

Precautions & Warnings

- Not recommended during pregnancy or breastfeeding
- Use with caution in patients with kidney or liver disorders
- Avoid alcohol to reduce the risk of side effects
- Inform your doctor about all existing medications to avoid interactions

Overdose Information

Do not exceed the prescribed dose. Overdose may cause serious complications including liver damage or breathing difficulty. In case of suspected overdose, seek immediate medical help.

Frequently Asked Questions

1. What is the primary use of OSVYNAC-MR Tablet?

To relieve musculoskeletal pain, reduce inflammation, and relax muscle stiffness caused by injury or overuse.

2. Can I stop taking the medication if I feel better?

No, always consult your doctor before discontinuing. Stopping early may cause symptoms to return.

3. Can it be used for rheumatoid arthritis?

It provides temporary relief from pain but does not treat the root cause of rheumatoid arthritis.

4. Is it safe for patients with kidney problems?

Only under medical supervision. Kidney function should be monitored.

5. Can it treat stomach pain?

No, and it may aggravate gastric symptoms. Do not use for abdominal pain.

6. How fast does it work?

Relief may begin within 30 minutes to 1 hour.

7. Can it be taken for a toothache?

It's not the ideal treatment. Consult your dentist for better options.

8. Is it safe to take with other painkillers?

No, unless directed by your doctor.

9. Can it cause liver damage?

Excessive use, particularly due to paracetamol, can harm the liver. Use only as directed.

10. What lifestyle changes help muscle recovery?

Stretching, gentle exercise, hydration, and proper rest can aid recovery while taking OSVYNAC-MR.

Move Freely. Live Comfortably.

OSVYNAC-MR Tablet supports your recovery by easing muscle pain and restoring mobility—so you can get back to your daily life with comfort and confidence.