

OSVYNAC-P Tablet

Dual Action Pain and Inflammation Relief

For joint, muscle, dental, and general body pain relief.

Product Overview

OSVYNAC-P Tablet is a fast-acting combination medication used to relieve pain, reduce inflammation, and manage fever. It is indicated for conditions such as rheumatoid arthritis, ankylosing spondylitis, osteoarthritis, muscle pain, back pain, dental pain, and discomfort in the ear or throat. Its formulation effectively targets the root causes of discomfort, helping improve daily function and quality of life.

Active Ingredients & Their Functions

Aceclofenac (100 mg)

- NSAID that blocks COX enzymes
- Reduces pain, inflammation, and swelling

Paracetamol (325 mg)

- Enhances pain relief and reduces fever
- Works synergistically with Aceclofenac for faster relief

Benefits of OSVYNAC-P Tablet

- Effective relief from joint and muscle pain
- Reduces swelling and improves mobility
- Alleviates fever and discomfort
- Suitable for short-term management of acute inflammatory pain

Recommended Uses

- Rheumatoid arthritis
- Osteoarthritis
- Ankylosing spondylitis
- Muscle, back, or dental pain
- Ear and throat discomfort

Directions for Use

- Take OSVYNAC-P with water after meals to reduce gastric irritation
- Do not chew, crush, or break the tablet
- Follow the prescribed dose and schedule

Storage Guidelines

- Store in a cool, dry place away from sunlight
- Keep out of reach of children

Side Effects

- Nausea or vomiting
- Dizziness
- Gastrointestinal disturbances (e.g., constipation, flatulence, diarrhea)
- Elevated liver enzymes (requires monitoring in long-term use)

How It Works

OSVYNAC-P Tablet combines Aceclofenac and Paracetamol to block pain signals and reduce inflammation. Aceclofenac inhibits the production of prostaglandins (pain-causing chemicals), while Paracetamol works on the brain's temperature regulation and pain perception centers.

Overdose Precautions

Do not exceed the recommended dose. An overdose may cause serious liver damage or other complications. Seek immediate medical help in case of accidental overdose.

Frequently Asked Questions (FAQs)

1. What is the main use of OSVYNAC-P Tablet?

It is used to manage pain, inflammation, and fever in various conditions.

2. Will it cure rheumatoid arthritis?

No, it provides symptomatic relief but is not a cure.

3. Can it be taken on an empty stomach?

No, it is best taken after food to reduce gastric irritation.

4. Can it be taken with antibiotics or vitamin supplements?

Consult your doctor before combining with other medications or supplements.

5. What if I forget a dose?

Take it as soon as remembered unless it's close to the next dose. Do not double the dose.

6. Can I stop taking it once I feel better?

Only under your doctor's supervision. Stopping early may cause relapse.

7. Is it suitable for children?

No, it is not recommended for pediatric use.

8. Can it be taken with other painkillers?

No, additional painkillers may increase the risk of side effects unless advised by a doctor.

9. How quickly does it work?

It usually starts working within 1.5 to 3 hours.

10. Can it cause dizziness or drowsiness?

Yes, avoid driving or using heavy machinery if affected.

Reliable Relief. Trusted Formula.

OSVYNAC-P Tablet is a dependable solution for relieving pain and inflammation, helping you return to your routine with comfort and ease.